

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K	A2	(11) International Publication Number: WO 97/13781 (43) International Publication Date: 17 April 1997 (17.04.97)
<p>(21) International Application Number: PCT/IL96/00119</p> <p>(22) International Filing Date: 26 September 1996 (26.09.96)</p> <p>(30) Priority Data: 115453 28 September 1995 (28.09.95) IL 118097 1 May 1996 (01.05.96) IL</p> <p>(71) Applicant (for all designated States except US): YEDA RESEARCH AND DEVELOPMENT CO. LTD. (IL/IL); Weizmann Institute of Science, P.O.B. 95, 76100 Rehovot (IL).</p> <p>(72) Inventors; and (75) Inventors/Applicants (for US only): CHEBATH, Judith (IL/IL); Rehov Miller 13, 76284 Rehovot (IL). HALIMI, Hubert (FR/FR); CNRS, IBMC, Laboratoire d'Immunochimie des Peptides et Virus, 15, rue des Cartes, F-67084 Strasbourg Cédex (FR). REVEL, Michel (IL/IL); Weizmann Institute of Science, Beit Brazil 5, 76100 Rehovot (IL).</p> <p>(74) Agent: EINAV, Henry; Inter-Lab Ltd., Science-based Industrial Park, Kiryat Weizmann, 76110 Ness-Ziona (IL).</p>		<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: SYNTHETIC PEPTIDES THAT INHIBIT IL-6 ACTIVITY</p> <p>(57) Abstract</p> <p>Novel peptides derived from the gp80 subunit of the IL-6 receptor system which inhibit the activity of IL-6 are provided.</p>		

<i>FOR THE PURPOSES OF INFORMATION ONLY</i>					
Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.					
AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia				

<i>FOR THE PURPOSES OF INFORMATION ONLY</i>					
Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.					
AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia				

<i>FOR THE PURPOSES OF INFORMATION ONLY</i>					
Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.					
AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia				

SYNTHETIC PEPTIDES THAT INHIBIT IL-6 ACTIVITY

Field of the Invention

The present invention is generally in the field of inhibitors of Interleukin-6 (IL-6) activity. More specifically, the present invention concerns new synthetic peptides which are capable of inhibiting the IL-6-dependent growth of myeloma/plasmacytoma cells.

Background of the Invention and Prior Art

The receptor system for Interleukin-6 (IL-6) is composed of two distinct receptor subunits designated gp80 (IL-6R) and gp130 (reviewed in Hirano et al., 1994). These two receptor proteins belong to the cytokine receptor superfamily of Bazan (1990). The 3-dimensional structure of the human Growth Hormone Receptor (hGHR), a member of this family, has been revealed by crystallography (DeVos et al., 1992), from which the residues which interact with the ligand and those which mediate the interaction between the two receptor subunits in their extracellular domains, have been determined. The alignment of the IL-6R and hGHR sequences based on Bazan's model is shown schematically in Figure 1.

IL-6 is a pleiotropic cytokine which has a number of important biological activities (see Revel, 1992, for review). Further, IL-6 has been implicated in the growth and progression of human multiple myeloma (Klein et al., 1990).

In fact, IL-6 is a growth factor for B-lymphocyte leukemic cells of Multiple Myeloma. These leukemic cells are also called plasmacytoma or myeloma cells, as they are derived from mature B-lymphocytes or plasma cells. When fused to antibody-producing B-cells, these myeloma or plasmacytoma cells are called hybridoma cells.

In view of the fact that IL-6 is a growth factor for such plasmacytoma or myeloma cells, there has been a long-felt need to obtain specific IL-6 inhibitors which may be used to block the IL-6-mediated or IL-6-dependent growth of such plasmacytoma/myeloma cells, and thereby provide a way for treating Multiple Myeloma, a disease affecting a very large number of people worldwide. To this end

peptide 249-264 (i.e. a peptide having the amino acid residues from residue No. 249 to residue No. 264 of the IL-6R amino acid sequence), which is capable of inhibiting the growth of murine plasmacytoma B9 cells. It is also known that IL-6 plays a role in the inhibition of other diseases, such as osteoporosis and autoimmune diseases.

5 Heretofore, neither the specific peptides of the present invention, nor their specific biological activities and other characteristics such as, for example, their specificity for certain monoclonal antibodies, have been described.

It is therefore an aim of the present invention to provide new peptides derived from IL-6R which are capable of inhibiting the IL-6-dependent growth of human
10 myeloma or murine plasmacytoma cells, osteoporosis and autoimmune diseases.

It is another aim of the present invention to provide such new peptides which are further characterized by virtue of their defining linear epitopes within the IL-6R sequence, which are the binding sites of monoclonal antibodies (Mabs) which are themselves capable of blocking IL-6 activity.

15 Yet another aim of the invention is to provide a chemical synthesis process for the preparation of the new peptides.

A yet further aim of the invention is to provide pharmaceutical compositions containing the new peptides.

20 Summary of the Invention

The present invention is based on the unexpected finding that short peptides within the IL-6 receptor gp80 molecule (IL-6R) could be defined by virtue of their ability to bind two different monoclonal antibodies (Mabs) which were previously known to strongly inhibit the activity of IL-6. Further, when chemically synthesized,
25 in accordance with the present invention, these peptides when added to cultures of leukemic cells, were surprisingly shown to be capable of causing the complete inhibition of the growth of such leukemic (plasmacytoma/myeloma) cells.

Accordingly, the present invention provides a peptide or biologically active analogs thereof capable of inhibiting the activity of IL-6, wherein said peptide is

system and by being a linear epitope recognized by one or more monoclonal antibodies (Mab) specific to IL-6R, with the proviso that said peptide is other than the group of peptides consisting of : (i) the 16 amino acid peptide having the amino acid sequence of residues 249-264 of the IL-6R molecule; (ii) the 14 amino acid peptide having the amino acid sequence of residues 255-268 of the IL-6R molecule; (iii) the 6 amino acid peptide having the amino acid sequence of residues 249-254 of the IL-6R molecule; (iv) the 10 amino acid peptide having the amino acid sequence of residues 259-268 of the IL-6R molecule; and (v) the 10 amino acid peptide having the amino acid sequence of residues 249-258 of the IL-6R molecule.

10 An embodiment of the above peptide or analogs thereof of the invention is a peptide selected from the group of peptides having between about 4 and about 25 amino acid residues derived from the portion of the IL-6R molecule extending between residue 223 and residue 272 as depicted in Fig. 2, said group of peptides comprising:

- 15 (a) a peptide having at least the 10 amino acid sequence LRYRAERSKT from position 255 to position 264 of the IL-6R molecule;
- (b) a peptide having at least the 10 amino acid sequence YRAERSKTFT from position 257 to position 266 of the IL-6R molecule;
- (c) a peptide having at least the 10 amino acid sequence AERSKTFTTW from position 259 to position 268 of the IL-6R molecule;
- 20 (d) a peptide having at least the 10 amino acid sequence RSKTFTTWMV from position 261 to position 270 of the IL-6R molecule;
- (e) a peptide having at least the 10 amino acid sequence KTFTTWMVKD from position 263 to position 272 of the IL-6R molecule;
- 25 (f) a peptide having at least the 10 amino acid sequence SFYRLRFELR from position 247 to position 256 of the IL-6R molecule;
- (g) a peptide having at least the 10 amino acid sequence YRLRFELRYR from position 249 to position 258 of the IL-6R molecule;
- (h) a peptide having at least the 10 amino acid sequence I RFFI RYRAF from position

(i) an analog of any one of the peptides of (a) - (h) in which one or more amino acid residues have been added, deleted or substituted by another amino acid residue, and wherein said peptides and analogs of (a) - (i) are characterized by defining all or part of a linear epitope recognized by anti-IL-6R monoclonal antibodies and by being
5 capable of inhibiting IL-6 activity.

Another embodiment of the above peptides or analogs of the invention is a peptide or analog wherein said peptide or analog defines a linear epitope recognized by one or both of the Mabs, herein designated Mab 34.4 and Mab 50.6.

A yet further embodiment of the above peptides of the invention is a peptide
10 selected from any one of the herein designated peptides: (i) 1062 having the sequence from position 247 to position 260 of the IL-6R molecule as depicted in Fig. 2; (ii) 1063 having the sequence from position 255 to position 270 of the IL-6R molecule as depicted in Fig. 2; (iii) 1086 having the sequence from position 226 to position 245 of the IL-6R molecule as depicted in Fig. 2; (iv) 1085 having the sequence from position
15 234 to position 245 of the IL-6R molecule as depicted in Fig. 2; and (v) 1122 having the sequence from position 260 to position 269 of the IL-6R molecule as depicted in Fig. 2.

The present invention also provides in another aspect, a peptide or biologically active analogs thereof capable of inhibiting IL-6 activity, wherein said peptide is
20 characterized by being derived from the gp80 (IL-6R) subunit of the IL-6 receptor system and by being part of all of the region of IL-6R that is involved in the binding to IL-6, with the proviso that said peptide is other than the group of peptides consisting of: (i) the 16 amino acid peptide having the amino acid sequence of residues 249-264 of the IL-6R molecule; (ii) the 14 amino acid peptide having the amino acid sequence
25 of residues 255-268 of the IL-6R molecule; (iii) the 6 amino acid peptide having the amino acid sequence of residues 249-254 of the IL-6R molecule; (iv) the 10 amino acid peptide having the amino acid sequence of residues 259-268 of the IL-6R molecule; and (v) the 10 amino acid peptide having the amino acid sequence of residues 249-258 of the IL-6R molecule

An embodiment of this aspect of the invention is a peptide selected from the herein designated peptides: (a) 983 having the sequence from position 277 to position 299 of the IL-6R molecule as depicted in Table 1; (b) 1064 having the sequence from position 290 to position 300 of the IL-6R molecule as depicted in Table 1; (c) 1067
5 having the sequence from position 125 to position 135 of the IL-6R molecule as depicted in Table 1; and (d) an analog of any one of (a) - (c) wherein one or more amino acids have been added, deleted or substituted by another amino acid.

The invention also provides a pharmaceutical composition comprising as active ingredient at least one of the above peptides, analogs or mixtures of any thereof and a
10 pharmaceutically acceptable carrier, diluent or excipient.

Embodiments of the pharmaceutical composition of the invention include a pharmaceutical composition for the inhibition of IL-6 and a pharmaceutical composition for the treatment of Multiple Myeloma, osteoporosis and autoimmune diseases.

15 In addition, the present invention also provides for the use of the above peptides, analogs or mixtures of any thereof for the preparation of any of the above pharmaceutical compositions, or for the inhibition of IL-6, or for the treatment of Multiple Myeloma, osteoporosis and autoimmune diseases.

Other aspects of the invention are set forth or arise directly from the following
20 detailed description of the invention.

Brief Description of the Drawings

Fig. 1 shows schematically a comparison of part of the amino acid sequence of the extracellular domain of the IL-6R molecule with that of the human Growth
25 Hormone Receptor (hGHR). The alignment of the sequences is according to Bazan (1990). The position of the epitopes for Mab 34.4 and 50.6 are shown by flags in the figure.

Fig. 2 shows schematically a segment of the IL-6R amino acid sequence in which the limits of the three peptides, designated herein 1086, 1062 and 1063, are

overlapping decapeptides with Mabs is depicted in the lower part of the figure, the decapeptides having sequences contained with the 1086, 1062 and 1063 peptides. In this lower part, the numbers refer to the position of each peptide on the membrane grid (spot-scan test), and the "+" symbols denote the relative intensity of the reactions (positive results), all as described in Examples 1-3.

Fig. 3 shows graphically the effects of peptides 1062, 1063 and control peptide 903 (see Table 1) on the growth of B9 plasmacytoma-derived cells which is dependent on IL-6 addition. Recombinant CHO-produced human IL-6 was used at 2.5 U/ml (i.e. 7.5 IU/ml) and the growth measured after 3 days. Peptides were added at indicated concentration together with IL-6. All as described in Examples 2 and 3

Fig. 4 shows graphically the effects of peptides 1062, 1063, 1067, 1086 and control peptide 903 (see Table 1) on growth of B9 plasmacytoma-derived cells, as described in Example 3.

Fig. 5 shows the sequences and the effects of peptides 1062, 1063, 1122, 1118 and control peptide 903 on growth of B9 plasmacytoma-derived cells, as described in Example 3.

Fig. 6 shows graphically a three-dimensional model of the IL-6 receptor complex comprising IL-6R, gp 130 and the IL-6 ligand. This model was computed on the basis of the crystallographic data of DeVos et al (1992) for hGHR. The position of peptide 1063 is shown by filled circles and the direction of the strands by empty arrows. Part of peptides 1062 is shown by open squares and peptides 1086 by open ellipses. All as also described in Example 3.

Detailed Description of the Invention

The present invention concerns in one aspect new peptides which are capable of inhibiting the activity of IL-6, which are characterized as being derived from the gp 80 (IL-6R) subunit of the IL-6 receptor system and as being linear epitopes recognized by one or more monoclonal antibodies (Mab) specific to IL-6R

IL-6 is a growth factor for B-lymphocyte leukemic cells of Multiple Myeloma, i.e. plasmacytoma or myeloma cells. In accordance with the present invention, the above new peptides, being short peptides within the IL-6R molecule were defined by their ability to bind two Mabs previously isolated and characterized (Novick et al., 1992) as being Mabs which strongly inhibited the activity of IL-6. These new peptides were synthesized chemically and added to cultures of such leukemic cells and were observed to cause the complete inhibition of the growth of such plasmacytoma/myeloma cells.

Thus, the new peptides or mixtures of two or more such peptides of the invention may be used in a general fashion for the inhibition of IL-6 and for the inhibition of IL-6-mediated cellular activities, when such activities are undesired. For example, when IL-6 is produced intracellularly in abnormally large amounts or when IL-6 is administered in large doses, and in both situations the IL-6 reaches undesirably high levels in the body, giving rise to undesirable side effects, the peptides or mixtures thereof according to the invention may be used. Alternatively, the new peptides individually or mixtures of two or more thereof may be used specifically for the treatment of diseases such as, for example, Multiple Myeloma in humans in which the growth of the cancerous cells constituting this disease (B-lymphocyte leukemic cells) is dependent of IL-6. In this application, the new peptides of the invention antagonize the binding of IL-6 to its receptor or interfere with the function of the receptor system which transduces, intracellularly, molecular signals leading to the growth of myeloma cells.

Of the above new peptides, preferred peptides are those having between about 4 and about 25 amino acid residues derived from the IL-6R molecule and which define linear epitopes recognized by either of the anti-IL-6R Mabs 34.4 and 50.6 (both Mabs being potent inhibitors of IL-6 activity). Of these preferred peptides are included : (a) a peptide having at least the 10 amino acid sequence LRYRAERSKT from position 255 to position 264 of the IL-6R molecule; (b) a peptide having at least the 10 amino acid sequence YRAFRSKTFT from position 257 to position 266 of the IL-6R

from position 259 to position 268 of the IL-6R molecule; (d) a peptide having at least the 10 amino acid sequence RSKTFTTWMV from position 261 to position 270 of the IL-6R molecule; (e) a peptide having at least the 10 amino acid sequence KTFTTWMVKD from position 263 to position 272 of the IL-6R molecule; (f) a peptide having at least the 10 amino acid sequence SFYRLRFELR from position 247 to position 256 of the IL-6R molecule; (g) a peptide having at least the 10 amino acid sequence YRLRFELRYR from position 249 to position 258 of the IL-6R molecule; (h) a peptide having at least the 10 amino acid sequence LRFELRYRAE from position 251 to position 260 of the IL-6R molecule.

Further, the above preferred peptides also include any of the herein designated peptides: (i) 1062 having the sequence from position 247 to position 260 of the IL-6R molecule as depicted in Fig. 2; (ii) 1063 having the sequence from position 255 to position 270 of the IL-6R molecule as depicted in Fig. 2; (iii) 1086 having the sequence from position 226 to position 245 of the IL-6R molecule as depicted in Fig. 2; (iv) 1085 having the sequence from position 234 to position 245 of the IL-6R molecule as depicted in Fig. 2, and (v) 1122 having the sequence from position 260 to position 269 of the IL-6R molecule as depicted in Fig. 2.

Moreover, it should be noted that in accordance with the present invention, the use of Mabs to define, isolate and characterize peptides which represent linear epitopes recognized by the Mabs on the IL-6R molecule, provides a new way in which to define and isolate peptides which are directly involved in the function of the IL-6R protein, for example peptides, such as those of the invention, which can interfere with the function of this protein, i.e. inhibit the activity of IL-6 (which is mediated by IL-6R). This use of Mabs is strikingly significant in view of the previous observations that the binding of Mabs to proteins is most often due to the recognition of a configuration within the protein and more rarely due to a recognition of a linear segment of the amino acid sequence of the protein.

In another aspect, the present invention also concerns new peptides which are capable of inhibiting the activity of IL-6 which are characterized by being derived

from the gp80 (IL-6R) subunit of the IL-6 receptor system and by being part or all of the region of the IL-6R that is involved in the binding to IL-6.

Thus, these other new peptides, while not being defined and isolated by virtue of their being linear epitopes recognized by known Mabs, nevertheless represent peptides which antagonize the binding of IL-6 to its receptor or interfere with the function of the receptor system which transduces, intracellularly, molecular signals leading to the growth of IL-6-dependent cells, e.g. myeloma cells. Accordingly, these peptides may be used in the same way as noted above, i.e. for the inhibition of IL-6 activity in general, when said inhibition is desired, or specifically for the inhibition of IL-6-dependent cancerous cells such as in Multiple Myeloma in humans.

Of these other new peptides, preferred peptides include: (a) 983 having the sequence from position 277 to position 299 of the IL-6R molecule as depicted in Table 1; (b) 1064 having the sequence from position 290 to position 300 of the IL-6R molecule as depicted in Table 1; (c) 1067 having the sequence from position 125 to position 135 of the IL-6R molecule as depicted in Table 1.

In yet another aspect, the present invention concerns biologically active analogs of any of the above noted new peptides. Suitable analogs are those which retain the ability to inhibit IL-6 activity by antagonizing the binding of IL-6 to its receptor or by interfering with the function of the receptor system which transduces, intracellularly, molecular signals leading to the growth of IL-6-dependent cells, such as, for example, myeloma cells. Of these suitable analogs are also included analogs which retain the essential amino acid residues of the linear epitopes recognized by the Mabs 34.4 and 50.6. These analogs of the invention are those which have at least one amino acid residue added, deleted or substituted by another amino acid when compared to the peptide amino acid sequence from which the analog amino acid sequence has been generated.

By virtue of the above characteristics of the analogs of the invention, these analogs may thus be used in the same way as the new peptides of the invention, i.e. as inhibitors of IL-6 activity in general or as agents for the treatment of IL-6-dependent

The peptides and the analogs of the invention may be prepared by any well known procedure of the art, in particular, by the well established chemical synthesis procedures utilizing automated peptide synthesizers followed by chromatographic purification (see Example 2).

5 In a further aspect, the invention also concerns pharmaceutical compositions containing as active ingredient any of the above new peptides, analogs or mixtures of two or more peptides or analogs, as well as any of the well known pharmaceutically acceptable carriers, diluents or excipients.

These pharmaceutical compositions may be formulated for any form of
10 administration as desired, for example, for intravenous, intraperitoneal or oral administration. Accordingly, the choice of diluent, carrier or excipient will be made according to the desired mode of administration.

The actual mode of administration and the optimal formulation therefor will be determined by skilled practitioners. Likewise, the dosage for administration and hence,
15 the concentration of the active ingredient in each dosage form according to its mode of administration, will also be decided by skilled practitioners. However, in this regard it is considered that a suitably effective dosage form, for any route of administration, will be one that will result in a concentration in the body of about 50-100 μ M of the peptides, analogs or mixtures thereof, following administration.

20 The pharmaceutical compositions of the invention may be used for the inhibition of IL-6 when such treatment is indicated, as noted above in cases where IL-6 levels are abnormally high, or they may be used specifically to treat such diseases where the diseases tissue is IL-6 dependent, for example, in Multiple Myeloma, in which the myeloma cells constituting the malignant growth are IL-6 dependent.

25 The present invention will now be described in more detail in the following non-limiting Examples, and the accompanying drawings.

Example 1: Epitopes recognized by Mab 34.4 and 50.6

Monoclonal antibodies 34.4 and 50.6, previously prepared and characterized by

IL-6 to the gp80 subunit of the IL-6 receptor (IL-6R) and consequently to inhibit IL-6 binding to human cells and its biological activity (Novick et al., 1992).

The two antibodies 34.4 and 50.6 were studied in accordance with the present invention as regards their ability to recognize linear peptides from the IL-6R extracellular domain. For this purpose, using standard chemical synthesis procedures (see also Example 2), a series of 96 decapeptides were synthesized which span residues 123-322 of the IL-6R sequence. Each consecutive peptide was offset by 2 amino acids from the previous one, yielding a nested, overlapping library (see IL-6R sequence in Fig. 1 and some of the peptides as shown in Fig. 2). A membrane carrying the 96 peptides was reacted with Mab 34.4 (18 μ g/ml) and stained with a β -galactosidase-conjugated secondary antibody (anti-mouse immunoglobulin antibody). Two series of β -galactosidase positive spots on the membrane were seen, one corresponding to residues 223-240 and the other to residues 241-272. A similar experiment was performed using a horseradish peroxidase-conjugated secondary antibody and detection by enhanced chemiluminescence (ECL kit, amersham Radiochemicals, UK). This method of detection is more quantitative and allowed for a comparison of the intensity of the reaction between the antibody and the diverse peptides. A strong reaction was observed with 4 peptide spots (67, 68, 69 and 70) corresponding to amino acids 255-264, 257-266, 259-268 and 261-270. These results are depicted in Fig. 2, which shows schematically a segment of the IL-6R amino acid sequence in which the limits of the three peptides designated herein as 1086, 1062 and 1063 are depicted by boxes (upper part of the figure) and which also shows the results of reaction of the various overlapping decapeptides contained within peptides 1086, 1062 and 1063 with Mabs (lower part of the figure). In these reaction results, the numbers refer to the position of each peptide on the membrane grid (spot-scan test) and the "+" symbols denote the extent of the reaction (+++ = very strong reaction; ++ = strong reaction; and + = fairly strong reaction). Thus, from these results, it appears

that the core of the epitope from Mab 34.4 is RSKT (261-264) but probably extends to

In a similar way, the 96-peptide library of IL-6R was also tested with Mab 50.6 with the ECL detection method. Spots 64 and 65 reacted most strongly indicating sequence 251-258 as the core epitope for this second neutralizing antibody (Fig. 2).

5 **Example 2: Synthetic peptides corresponding to epitopes defined by Mab 34.4 and 50.6 inhibit the biological activity of IL-6 on hybridoma cell growth**

The region of the IL-6R sequence to which the neutralizing Mab 34.4 and 50.6 bind was determined as noted above (Fig. 2). The indicated peptides 1062 (IL-6R residues 257-260) 1063 (255-270) and 1086 (226 and 245) were obtained by solid-
10 phase synthesis with a peptides synthesizer from Applied Biosystems (Foster City, CA). Purity of the peptides was verified by Reverse Phase High Performance Liquid Chromatography (RP-HPLC) on an Aquapore C8 column (Brownlee).

These purified peptides were then tested for their ability to interfere with IL-6 action on cells. The rationale was that since the antibodies blocking IL-6 activity bind
15 to epitope sequences within these peptides, it is likely that these sequences are important for some protein-protein interactions within the receptor complex. Hence, these peptides could themselves mimic such interactions and interfere with the normal protein-protein contacts necessary for the function of the receptor system.

To assay the biological activity of the peptides as inhibitors of IL-6 action
20 against growth of plasmacytoma cells, a hybridoma cell line B9 whose growth is IL-6 dependent (Helle et al, 1988) was used. The B9 cells were cultured in suspension in growth medium RPMI 1640 (Biolabs Ltd., Israel) with 10% heat-inactivated fetal calf serum and 4 U/ml IL-6. The recombinant human IL-6 from Chinese Hamster Ovary (CHO) cells was a pure preparation as described (Novick et al., 1989) which titrated
25 on T1165 plasmacytoma cells 5×10^6 units/mg (corresponding to 1.5×10^7 International units/mg). Units used here are T1165 units (1 unit = 3 International units).

Prior to assay, the B9 cells were washed and incubated for 5 hours in growth medium but without IL-6. The cells were centrifuged, resuspended in growth medium with 2.5 units/ml (i.e. 7.5 IU/ml) of IL-6 and seeded in a 96-well microplate (Nunc,

solution with appropriate peptide concentration was included in the final volume of each well, i.e. the peptides at concentrations ranging from 62.5 $\mu\text{g/ml}$ to 250 $\mu\text{g/ml}$ (as well as a 0 $\mu\text{g/ml}$ control) were added together with the IL-6. After 3 days of culture at 37°C, a solution of Alamar Blue (Biosource, Camarillo, CA) was added at 20 $\mu\text{g/ml}$ per well and after mixing, incubation was continued for 3 hours. The optical absorbance at 530 nm wavelength of each well was measured as specified by the manufacturer. The absorbance is directly proportional to the number of living cells in the well as verified in calibration experiments from 50,000 to 10^6 B9 cells (0.25 OD530 corresponding to 5×10^5 B9 cells under the present experimental conditions). The results were expressed as the difference between final cell density with IL-6 and without IL-6. The IL-6 dependent difference in the absence of peptide was defined as 100% growth.

The results of the above experiments are given in Fig. 4, which shows graphically the effects of peptides 1062, 1063 and control peptide (see below and Table 1) on the IL-6-dependent growth of the B9 plasmacytoma-derived cells. As is apparent from these results, the addition of increasing concentrations of peptide 1063 (255-270) resulted in a dose dependent inhibition of the growth of the B9 cells. At the highest dose, the growth inhibition was 99%. A complete inhibition was also obtained with peptide 1062 (257-260) but with a different dose curve. In contrast, a control peptide, 903 (with an unrelated sequence as shown in Table 1 below), did not inhibit the IL-6 dependent growth of the B9 cells. Therefore, the two peptides 1062 and 1063 corresponding to the core epitopes of the neutralizing Mabs 34.4 and 50.6 are potent specific inhibitors of the IL-6-dependent growth of plasmacytoma-derived cells.

TABLE 1

Peptides tested for inhibition of IL-6 activity and their coordinates in IL-6R

Peptide	Sequence	Coordinates	Remarks
1062	SFYRLRFELRYRAE	247-260	Mab 50.6 epitope
1063	LRYRAERSKTFTTWMV	255-270	Mab 34.4 epitope
983	CVIHDAWSGLRHVVQLRAQEEFG	277-299	
1064	VQLRAQEEFGQ	290-300	IL-6 binding
1067	SPLSNVCEWG	125-135	N-domain start
1085	LSVTWQDPHSWN	234-245	
1086	AVARNPRWLSVTWQDPHSWN	226-245	gp130 interaction?
903	PGHRYRDQQTQTSFSEEPQSSQLLPC	other protein	control peptide

5

The activity of peptide 1063 (255-270, Mab 34.4 epitope) was tested in 5 separate experiments and the inhibition at the three doses was statistically highly significant (Table 2 below). Direct comparison of peptide 1063 to control peptide 903 showed a highly significant inhibition of the IL-6 dependent growth of the plasmacytoma derived B9 cells even when peptide 1063 was used at half the concentration of the control peptide (see legend of Table 2 below).

15

20

TABLE 2**IL-6 Antagonist activity of peptide from Mab 34.4 and 50.6 epitopes**

PEPTIDE	IL-6 DEPENDENT HYBRIDOMA B9 CELL GROWTH		
	62.5 µg/ml	125 µg/ml	250 µg/ml
	Percent of IL-6 dependent cell growth ± SD		
1063 (255-270)	71±9 (p=0.001)	45±16 (p<0.001)	2±1 (p<0.0001)
1062 (257-260)	100±0 (n.s.)	15±9 (p<0.0001)	
903 (control)	97±4 (n.s.)	95±9 (n.s.)	89±15 (n.s.)

- 5 Statistical analysis of 5 experiments using two-tailed Student's t-test in comparison to no peptide. comparison of B9 cell growth with peptide 1063 and 125 µg/ml to control peptide 903 at 250 µg/ml was highly significant at p=0.004.

10 **Example 3: Comparison of peptides from Mab 34.4 and 50.6 epitopes to peptides from other regions of the IL-6R sequence for inhibition of hybridoma B9 cell growth.**

Five peptides were prepared as noted above (Example 2) corresponding to different regions of the IL-6R sequence (see Table 1 above). Peptides 983 and 1064 correspond to a region homologous to that which in hGHR is involved in ligand binding (Fig. 1; DeVos et al., 1992) and which contains the RAQEEF motif shown to
 15 bind IL-6 (Martin et al., 1994). Peptide 1067 also corresponds to a region which in the hGHR molecule is involved in ligand binding (Fig. 1). Peptide 1086 corresponds to a region which has been shown to interact with gp130 (Savino et al., 1994) and is close to the epitope of Mab 34.4 and 50.6 (Fig. 2). These peptides were tested for their
 20 effect on the IL-6 dependent growth of the plasmacytoma-derived B9 cells, as in Example 2.

A comparison of the residual IL-6 dependent growth (see Table 3 below) shows that overlapping peptides 983 and 1064 were not active as compared to peptide 1063 (255-270). Peptide 1067 did show some activity, but lower than peptide 1063. In a

62% \pm 43 and was not significant statistically. However, peptide 1086 (226-245) did show activity, although lower than that of peptides 1062 and 1063. Fig. 4 shows graphically the effects of peptides 1062, 1063, 1067 and 1086 as compared to each other and to control peptide 903, on the growth of B9 plasmacytoma-derived cells.

5 The results shown are the mean results of 4 experiments.

Fig. 5 shows again the sequences of the above mentioned peptides 1062 and 1063, and their effect on the growth of B9 plasmacytoma-derived cells, as well as those of peptides 1118 and 1122, having low, and strong effect, respectively, and those of two related peptides found to be inactive.

10

TABLE 3

Comparison of different peptides

	HYBRIDOMA B9 CELL GROWTH WITH PEPTIDE				
	1063	1064	983	1067	903
	Percent of IL-6 dependent cell growth				
IL-6 : 1 U/ml	3%	74%	100%	48%	100%

15 Peptides tested at 250 μ g/ml

A three-dimensional model of the IL-6R molecule was produced based on the crystallographic coordinates of hGHR. This is shown schematically in Fig. 5, which is a three-dimensional model of the IL-6R complex comprising IL-6R, gp130 and the IL-6 ligand. This model was computed on the basis of the crystallographic data of DeVos (1992) for hGHR. The position of the different peptides tested is indicated. Peptide 1063 (255-270, marked by filled circles) does not appear to interact with the IL-6 ligand. Peptide 1062 (247-260, marked by open squares in the region preceding its overlap with peptide 1063) does contain residues around position 247 which could interact with IL-6. Peptide 1086 (226-245, marked by open ellipses) contains a loop

20

25

IL-6R binding (Hirano et al., 1994) and the IL-6R may similarly dimerize to form a hexameric complex. Building a computer model of this hexameric complex by a symmetric assembly of two of the trimeric complexes shown in Fig. 5, revealed that peptide 1063 and part of 1062 are engaged in a second interaction of IL-6R and gp130. Therefore, the regions common to the active peptides, in particular peptide 1063 which is the most active inhibitor of IL-6, appear to interfere with the interactions between IL-6R and gp130 to form the hexameric complex, and hence inhibit the function of the IL-6 receptor complex.

10 **Example 4: Peptides defined by the Mab 34.4 and 50.6 epitopes inhibit the IL-6 dependent growth of human myeloma cells**

Multiple myeloma is a leukemia of plasma cells in which IL-6 is considered as a promoting growth factor (Klein et al., 1990). It was, therefore, of interest to determine if the peptides derived from the epitopes Mab 34.4 and 50.6, or other peptides, would inhibit the growth of such tumor cells. The XG-1 cell line has been established from a patient with multiple myeloma (Klein et al., 1990) and the growth of these cells is dependent on the addition of IL-6. Cultures of XG-1 cells were established in 6-well Costar plates, in a final volume of 1 ml. The cells were cultured with or without IL-6 (0.2 U/ml, i.e. 0.6 IU/ml) for 3 days and then pulsed with tritium-labeled thymidine for 2 hours. The incorporation of radioactive thymidine was used as a measure of DNA replication and growth. As indicated in Table 4 below, the IL-6 dependent growth was strongly reduced (by 85%) by the addition of 100 µg/ml of peptide 1062, and was reduced by more than 55% with peptide 1063. In contrast peptide 1064, gave only a small inhibitory effect. Peptide 1067 was found also to be active to inhibit the growth of these human myeloma cells (see Table 4 below).

Therefore, the IL-6R peptides are capable of inhibiting the function of the human IL-6 receptor complex in human tumor cells.

TABLE 4**Effect on human myeloma XG-1 cells**

	HUMAN MYELOMA XG-1 GROWTH WITH PEPTIDE				
	none	1062	1063	1064	1067
	³ H-Thymidine incorporation, cpm x 10 ⁻³				
No IL-6	6.4±1.6	8.3±0.3	8.7±2.5	10.1±0.3	6.5±1.9
IL-6, 0.2 U/ml	43.6±0.9	15.1±3.3	25.2±0.2	39.1±1.5	10.7±0.3
IL-6 dpdt (%)	37.2 (100)	6.8 (18.2)	16.5 (44.3)	29.0 (78)	4.2 (11.3)

5

In view of the results set forth herein above (Examples 1-4), a comparison was made between these results ("Present Invention") and the previously reported results concerning peptides described by Cochrane (Grube and Cochrane, 1994, "Prior Art"). This comparative analysis is set forth in Table 5 below. It should be noted that

10 Cochrane showed only peptide 249-264 (Y249-T264) to be active on the B9 lymphocytoma-derived cells, whilst the other peptides described by Cochrane were tested on HepG2 cell for effects on fibrinogen secretion. In contrast, the peptides of the present invention 1062 (247-260), 1063 (255-270), 1086 (226-245), and 1122 (260-269) which all differ (although with some overlap as regards 1062 and 1063)

15 from those of Cochrane, have been shown to be active on B9 cells. Moreover, peptides 1062 and 1063 inhibit human myeloma XG-1 cells. Furthermore, the most similar peptides (although still being different) of the present invention and of Cochrane, namely 1063 (255-270) and 255-268, respectively, are also different in activity. The peptide of the present invention, 1063 (L255-V270) has been shown in accordance

20 with the invention to be active on B9 cells whilst that of Cochrane (255-268) tested on HepG2 cells was inactive.

Finally, the peptides prepared and characterized according to the present invention have been isolated by the initial procedure of epitope mapping using the

previously prepared Mabs 34.4 and 50.6. In contrast, Cochrane prepared the peptides by random synthesis of different peptides.

TABLE 5

5 **Comparison between the activity of the peptides of the present invention
and those of the prior art (Grube and Cochrane, 1994)**

Present Invention			Prior Art		
PEPTIDE	RESULT		PEPTIDE	RESULT	
	B9 cells	XG-1		B9 cells	HepG2
247-260 (1062)	active	active	249-264	active	active
255-270 (1063)	active	somewhat	255-268	not done	inactive
			249-254	not done	inactive
			259-268	not done	inactive
			249-258	not done	active
226-245 (1086)	active	not done			

References

- Bazan J.F. (1990) Structural design and molecular evolution of a cytokine receptor superfamily. *Proc. Natl. Acad. Sci. USA* 87:6934-6938.
- 5 DeVos A.M., Ultsch M. and Kossiakoff A.A. (1992) Human Growth Hormone and extracellular domain of its receptor: crystal structure of the complex. *Science* 255: 306-312.
- 10 Grube B.J. and Cochrane C.G. (1994), Identification of a regulatory domain of the Interleukin-6 receptor. *J. Biol. Chem.* 269:20791-20797.
- Helle M., Boije L. and Aarden L. (1988) Functional discrimination between Interleukin-6 and Interleukin-1. *Eur. J. Immunol.* 18:1535-1540.
- 15 Hirano T., Matsuda T. and Nakajima K. (1994) Signal transduction through gp130 that is shared among the receptors for the Interleukin-6 related cytokine subfamily. *Stem Cells* 12:262-277.
- 20 Klein B., Zhang X.G., Jourdan M., Boiron J.M., Portier M., Lu Z.Y., Wijdenes J., Brochier J. and Bataille R. (1990) Interleukin-6 is the central tumor growth factor in vitro and in vivo in multiple myeloma. *Eur. Cyt. Netw.* 1: 193-201.
- Martin F., Toniatti C., Salvati A.L., Venturini S., Ciliberto G., Cortese R. and
25 Sollazzo M. (1994) The affinity selection of a minibody polypeptide inhibitor of human IL-6. *EMBO J.* 13:5303-5309.
- Novick D., Eshhar Z., Revel M. and Mory Y. (1989) Monoclonal antibodies for affinity purification of IL-6/IFN- β 2 and for neutralization of HGF activity.

Novick D., Shulman L.M., Chen L. and Revel M. (1992) Enhancement of Interleukin-6 cytostatic effect on human breast carcinoma cells by soluble IL-6 receptor from urine and reversion by Monoclonal Antibody. *Cytokine* 4:6-11.

5

Revel M. (Ed.) IL-6: Physiopathology and clinical potentials. Sero Symposia Publications, Vol. 88, Raven Press, New York, 1992.

Savino R., Ciapponi L., Lahm A., Demartis A., Cabibbo A., Toniatti C., Delmastro P.,
10 Altmura S. and Ciliberto G. (1994) Rational design of a receptor superantagonist human IL-6. *EMBO J.* 13:5863-5870.

Claims

1. A peptide or biologically active analogs thereof capable of inhibiting the activity of IL-6, wherein said peptide is characterized by being derived from the gp80 (IL-6R) subunit of the IL-6 receptor system and by being a linear epitope recognized by one or more monoclonal antibodies (Mab) specific to IL-6R, with the proviso that said peptide is other than the group of peptides consisting of : (i) the 16 amino acid peptide having the amino acid sequence of residues 249-264 of the IL-6R molecule; (ii) the 14 amino acid peptide having the amino acid sequence of residues 255-268 of the IL-6R molecule; (iii) the 6 amino acid peptide having the amino acid sequence of residues 249-254 of the IL-6R molecule; (iv) the 10 amino acid peptide having the amino acid sequence of residues 259-268 of the IL-6R molecule; and (v) the 10 amino acid peptide having the amino acid sequence of residues 249-258 of the IL-6R molecule.
2. A peptide or analogs thereof, according to claim 1, selected from the group of peptides having between about 4 and about 25 amino acid residues derived from the portion of the IL-6R molecule extending between residue 223 and residue 272 as depicted in Fig. 2, said group of peptides comprising:
- (a) a peptide having at least the 10 amino acid sequence LRYRAERSKT from position 255 to position 264 of the IL-6R molecule;
 - (b) a peptide having at least the 10 amino acid sequence YRAERSKTFT from position 257 to position 266 of the IL-6R molecule;
 - (c) a peptide having at least the 10 amino acid sequence AERSKTFTTW from position 259 to position 268 of the IL-6R molecule;
 - (d) a peptide having at least the 10 amino acid sequence RSKTFTTWMV from position 261 to position 270 of the IL-6R molecule;
 - (e) a peptide having at least the 10 amino acid sequence KTFTTWMVKD from position 263 to position 272 of the IL-6R molecule;

- (f) a peptide having at least the 10 amino acid sequence SFYRLRFELR from position 247 to position 256 of the IL-6R molecule;
- (g) a peptide having at least the 10 amino acid sequence YRLRFELRYR from position 249 to position 258 of the IL-6R molecule;
- 5 (h) a peptide having at least the 10 amino acid sequence LRFELRYRAE from: position 251 to position 260 of the IL-6R molecule; and
- (i) an analog of any one of the peptides of (a) - (h) in which one or more amino acid residues have been added, deleted or substituted by another amino acid residue, and wherein said peptides and analogs of (a) - (i) are characterized by
- 10 defining all or part of a linear epitope recognized by anti-IL-6R monoclonal antibodies and by being capable of inhibiting IL-6 activity.
3. A peptide or analog thereof according to claim 1 or claim 2, wherein said peptide or analog defines a linear epitope recognized by one or both of the Mabs, herein
- 15 designated Mab 34.4 and Mab 50.6.
4. A peptide according to any one of claims 1-3, wherein said peptide is selected from any one of the herein designated peptides: (i) 1062 having the sequence from position 247 to position 260 of the IL-6R molecule as depicted in Fig. 2; (ii)
- 20 1063 having the sequence from position 255 to position 270 of the IL-6R molecule as depicted in Fig. 2; (iii) 1086 having the sequence from position 226 to position 245 of the IL-6R molecule as depicted in Fig. 2; and (iv) 1085 having the sequence from position 234 to position 245 of the IL-6R molecule as depicted in Fig. 2, and 1122 having the sequence from position 260 to position 269 of the
- 25 IL-6R molecule as depicted in Fig. 2.
5. A peptide or biologically active analogs thereof capable of inhibiting IL-6 activity, wherein said peptide is characterized by being derived from the gp80 (II -6R) subunit of the II -6 receptor system and by being part of all of the region

is other than the group of peptides consisting of : (i) the 16 amino acid peptide having the amino acid sequence of residues 249-264 of the IL-6R molecule; (ii) the 14 amino acid peptide having the amino acid sequence of residues 255-268 of the IL-6R molecule; (iii) the 6 amino acid peptide having the amino acid sequence of residues 249-254 of the IL-6R molecule; (iv) the 10 amino acid peptide having the amino acid sequence of residues 259-268 of the IL-6R molecule; and (v) the 10 amino acid peptide having the amino acid sequence of residues 249-258 of the IL-6R molecule.

6. A peptide or analog according to claim 5, wherein said peptide is selected from the herein designated peptides: (a) 983 having the sequence from position 277 to position 299 of the IL-6R molecule as depicted in Table 1; (b) 1064 having the sequence from position 290 to position 300 of the IL-6R molecule as depicted in Table 1; (c) 1067 having the sequence from position 125 to position 135 of the IL-6R molecule as depicted in Table 1; and (d) an analog of any one of (a) - (c) wherein one or more amino acids have been added, deleted or substituted by another amino acid.

7. A pharmaceutical composition comprising as active ingredient at least one peptide or analog thereof or mixtures of two or more peptides or analogs thereof according to any one of claims 1-4, and a pharmaceutically acceptable carrier, diluent or excipient.

8. A pharmaceutical composition according to claim 7 for the inhibition of IL-6.

9. A pharmaceutical composition according to claim 7 or claim 8 for the treatment of Multiple Myeloma.

10. A pharmaceutical composition according to any one of claims 7-9, wherein the

is sufficient to provide a 50-100 μ M concentration in the body following administration of said pharmaceutical composition.

- 5 11. Use of a peptide or analog thereof or mixtures of two or more peptides or analogs thereof according to any one of claims 1-4 for the preparation of a pharmaceutical composition.
- 10 12. Use of a peptide or analog thereof or mixtures of two or more peptides or analogs thereof according to any one of claims 1-4 for the preparation of a pharmaceutical composition for the treatment of Multiple Myeloma, osteoporosis, and autoimmune diseases.
- 15 13. Use of a peptide or analog thereof or mixtures of two or more peptides or analogs thereof according to any one of claims 1-4 for the inhibition of IL-6 activity.
14. Use of a peptide or analog thereof or mixtures of two or more peptides or analogs thereof according to any one of claims 1-4 for the treatment of Multiple Myeloma.
- 20 15. A peptide, analog, or mixtures of peptides or analogs thereof according to any one of claims 1-4 for use in the inhibition of IL-6.
16. A peptide, analog, or mixtures of peptides or analogs thereof according to any one of claims 1-4 for use in the treatment of Multiple Myeloma.
- 25 17. A peptide, analog, or mixtures of peptides or analogs thereof according to any one of claims 1-4 for use in the preparation of a pharmaceutical composition.
18. A pharmaceutical composition comprising as active ingredient at least one

according to claim 5 or claim 6, and a pharmaceutically acceptable carrier, diluent or excipient.

19. A pharmaceutical composition according to claim 18 for the inhibition of IL-6.

5

20. A pharmaceutical composition according to claim 18 or claim 19 for the treatment of Multiple Myeloma.

21. Use of a peptide, analog, or mixtures or peptides or analogs according to claim 5 or claim 6 for the preparation of a pharmaceutical composition.

10

22. A peptide, analog, or mixtures or peptides or analogs according to claim 5 or claim 6 for use in the treatment of Multiple Myeloma.

15

20

25

1/6

ALIGNEMENTS IL-6R and HGHR

Bold residues for ligand binding
 Italic residues for HGHR dimerization
 Circles: mutation impairing IL-6 binding


Epitopes of McABs

	121	oo	134	o	144	151
IL-6R	PPEEPQLSCFRKS	P	LSNVVCEWGP	RSTPSL		TTKAVLLVRKFQN
HGHR	SKEPKFTKCRSPE		RETFSCHWTDEVH	HGTK	*	LGPIQLFYTRRNT
	25	38	43	48		70
	160	170	180	oo	190	oo
IL-6R	SP	AEDFQEP	CQYSQESQ	KFSCQLAVPEG	DSSFYIVSMCVASS	
HGHR	**	WTQEWKECP	DYVSAG	ENSCYFNSSFTS	IWIPYCIKLT	SNGG
			218	ooo		
IL-6R		VGSKFSKTQT	FQCGIL	QPDPPA		
HGHR		TVDEKCF	SVDEIV	QPDPPI		
					McAB50.6	McAB34.4
	221	226	233	o	238	o
IL-6R	NITVTAVA		RN		PRWLSVTWQDP	HSWNSS
HGHR	ALNWTLLN	*	SLT**		HADIQVRWEAP	RNADIQ
					K*	WMVLEYELQYKEVNE
			165	169		
	267	270	277	oo	282	287
IL-6R	KTFTTWMV	KDLQH		HCVIHDAWS	GLRHVVQLRAQEEF	GQGEWSEWS
HGHR	TKWKMDP	PILTT		SVPVYSLKV	DKEYEVRVR	SKQR
					NSGNYGEFS	
	309	314				
IL-6R	PEAMGT	PWTESRSPP				
HGHR	EVLYVTLP	QMSQFT				

Figure 1

2/6

FIG. 2 : EPITOPES OF McAB 34.4 AND 50.6 IN IL-6 RECEPTOR gp80



PEPTIDES RECOGNIZED ON SPOT-SCAN

McAB 34.4	67	LYRAERSKT	++
	68	YRAERSKTFT	+++
	69	AERSKTFTTW	++
	70	RSKTFTTWVM	++
	71	KTFTTWVMVKD	+
McAB 50.6	63	SFYRLRFELR	+
	64	YRLRFELRYR	++
	65	LRFELRYRAE	+++

Figure 2

3/6


INHIBITION OF B9 HYBRIDOMA GROWTH BY IL-6R PEPTIDES

Figure 3


INHIBITION OF B9 HYBRIDOMA GROWTH BY IL-6R PEPTIDES

Figure 4

5/6

EFFECT OF IL-6R PEPTIDES
ON GROWTH OF PLASMACYTOMA B9 CELLS

247 255 260 265 270 275 280 285
 SFYRLRFELRYRAERSKTFTTWMVKDLQHHCVIHDAWSG
 β strand C Loop C-C' β strand C' β strand E

SFYRLRFELRYRAE
 LRYRAERSKTFTTWMV

1062 Act

1063 Act

YRLRFE

inactive


1122 Act

ERSKTFTTWM
 AERSKTFTTW

inactive

1118 Low

QHHCVIHDAWSG






Figure 6